Author Index

Ackermans, M. T., see Beckers, J. L. 537(1991)407

Ackermans, M. T., see Beckers, J. L. 537(1991)429

Akiyama, T., see Kuno, Y. 537(1991)489

Al-Ammar, A. S., Hamid, H. A., Rashid, B. H. and Basheer, H. M.
Preseparation of uranium by reversed-phase partition chromatography. Application to the determination of trace elements in nuclear-grade uranium compounds by inductively coupled plasma atomic emission spectrometry 537(1991)287

Alexandrovich, O., see Sigalov, A. 537(1991)464

Ang, H. G., Kwik, W. L. and Leong, W. K.
High-performance liquid chromatography of substituted trinuclear osmium carbonyl clusters, $Os_3(CO)_{12-n}[P(C_6F_5)_3]_n$ ($n = 0$, 1, 2) 537(1991)475

Basheer, H. M., see Al-Ammar, A. S. 537(1991)287

Beckers, J. L., Everaerts, F. M. and Ackermans, M. T.
Determination of absolute mobilities, pK values and separation numbers by capillary zone electrophoresis. Effective mobility as a parameter for screening 537(1991)407

Beckers, J. L., Everaerts, F. M. and Ackermans, M. T.
Isotachophoresis with electroosmotic flow: open *versus* closed systems 537(1991)429

Bezuidenhout, H. S., see Jones, E. A. 537(1991)277

Bourque, A. J. and Krull, I. S.
Solid-phase reagent containing the 3,5-dinitrophenyl tag for the improved derivatization of chiral and achiral amines, amino alcohols and amino acids in high-performance liquid chromatography with ultraviolet detection 537(1991)123

Bourrier, M.-J., see Goiffon, J.-P. 537(1991)101

Boyer, G. L., see Speirs, R. J. 537(1991)259

Brun, M., see Goiffon, J.-P. 537(1991)101

Byerley, T. J., see Fanska, C. B. 537(1991)357

Carini, M., see Pietta, P. 537(1991)449

Cartoni, G. P., Coccioli, F., Pontelli, L. and Quattrucci, E.
Separation and identification of free phenolic acids in wines by high-performance liquid chromatography 537(1991)93

Castello, G., see Gerbino, T. C. 537(1991)305

Chang, F. C., see Iida, T. 537(1991)345

Coccioli, F., see Cartoni, G. P. 537(1991)93

Cox, G. B., see Snyder, L. R. 537(1991)507

Crimmins, D. L., see Thoma, R. S. 537(1991)153

Czok, M. and Guiochon, G.
Pitfalls in the choice of isotherms for the calculation of band profiles in preparative chromatography 537(1991)497

Devani, M. B., Shishoo, C. J., Shah, S. A., Soni, K. P. and Shah, R. S.
Localization of amino acids on thin-layer chromatograms with acetylacetone–formaldehyde reagent 537(1991)494

Diwan, A. M., see Elias, C. 537(1991)460

Eick, J. D., see Fanska, C. B. 537(1991)357

Elias, C. and Diwan, A. M.
Behaviour of concanavalin A on DEAE-Sephadex and CM-Sephadex 537(1991)460

Ellegren, H., see Kågedal, L. 537(1991)17

Engström, B., see Kågedal, L. 537(1991)17

Everaerts, F. M., see Beckers, J. L. 537(1991)407

Everaerts, F. M., see Beckers, J. L. 537(1991)429

Facino, R. M., see Pietta, P. 537(1991)449

Fanska, C. B., Byerley, T. J. and Eick, J. D.
Indirect determination of isocyanates by gas chromatography 537(1991)357

Finn, N., see Smith, R. M. 537(1991)51

Ford, S. H., Gallery, J., Nichols, A. and Shambee, M.
High-performance liquid chromatographic analysis of the (cyanoaquo) stereoisomers of several putative vitamin B_{12} precursors 537(1991)235

Furusawa, M., see Kiba, N. 537(1991)443

Gallery, J., see Ford, S. H. 537(1991)235

Gardana, C., see Pietta, P. 537(1991)449

Gerbino, T. C. and Castello, G.
Gas chromatographic identification of complex mixtures of halomethanes and haloethanes by using the correlation between their retention and vapour pressure 537(1991)305

Goiffon, J.-P., Brun, M. and Bourrier, M.-J.
High-performance liquid chromatography of red fruit anthocyanins 537(1991)101

Goto, J., see Iida, T. 537(1991)345

Goyal, S. S., Hafez, A. and Rains, D. W.
Simultaneous determination of arsenite, arsenate, selenite and selenate in waters using suppressed ion chromatography with ultraviolet absorbance detection 537(1991)269

Guiochon, G., see Czok, M. 537(1991)497

Gülz, P.-G., see Vogt, T. 537(1991)453

Author Index

Ackermans, M. T., see Beckers, J. L. 537(1991)407

Ackermans, M. T., see Beckers, J. L. 537(1991)429

Akiyama, T., see Kuno, Y. 537(1991)489

Al-Ammar, A. S., Hamid, H. A., Rashid, B. H. and Basheer, H. M.
Preseparation of uranium by reversed-phase partition chromatography. Application to the determination of trace elements in nuclear-grade uranium compounds by inductively coupled plasma atomic emission spectrometry 537(1991)287

Alexandrovich, O., see Sigalov, A. 537(1991)464

Ang, H. G., Kwik, W. L. and Leong, W. K.
High-performance liquid chromatography of substituted trinuclear osmium carbonyl clusters, $Os_3(CO)_{12-n}[P(C_6F_5)_3]_n$ ($n = 0$, 1, 2) 537(1991)475

Basheer, H. M., see Al-Ammar, A. S. 537(1991)287

Beckers, J. L., Everaerts, F. M. and Ackermans, M. T.
Determination of absolute mobilities, pK values and separation numbers by capillary zone electrophoresis. Effective mobility as a parameter for screening 537(1991)407

Beckers, J. L., Everaerts, F. M. and Ackermans, M. T.
Isotachophoresis with electroosmotic flow: open *versus* closed systems 537(1991)429

Bezuidenhout, H. S., see Jones, E. A. 537(1991)277

Bourque, A. J. and Krull, I. S.
Solid-phase reagent containing the 3,5-dinitrophenyl tag for the improved derivatization of chiral and achiral amines, amino alcohols and amino acids in high-performance liquid chromatography with ultraviolet detection 537(1991)123

Bourrier, M.-J., see Goiffon, J.-P. 537(1991)101

Boyer, G. L., see Speirs, R. J. 537(1991)259

Brun, M., see Goiffon, J.-P. 537(1991)101

Byerley, T. J., see Fanska, C. B. 537(1991)357

Carini, M., see Pietta, P. 537(1991)449

Cartoni, G. P., Coccioli, F., Pontelli, L. and Quattrucci, E.
Separation and identification of free phenolic acids in wines by high-performance liquid chromatography 537(1991)93

Castello, G., see Gerbino, T. C. 537(1991)305

Chang, F. C., see Iida, T. 537(1991)345

Coccioli, F., see Cartoni, G. P. 537(1991)93

Cox, G. B., see Snyder, L. R. 537(1991)507

Crimmins, D. L., see Thoma, R. S. 537(1991)153

Czok, M. and Guiochon, G.
Pitfalls in the choice of isotherms for the calculation of band profiles in preparative chromatography 537(1991)497

Devani, M. B., Shishoo, C. J., Shah, S. A., Soni, K. P. and Shah, R. S.
Localization of amino acids on thin-layer chromatograms with acetylacetone–formaldehyde reagent 537(1991)494

Diwan, A. M., see Elias, C. 537(1991)460

Eick, J. D., see Fanska, C. B. 537(1991)357

Elias, C. and Diwan, A. M.
Behaviour of concanavalin A on DEAE-Sephadex and CM-Sephadex 537(1991)460

Ellegren, H., see Kågedal, L. 537(1991)17

Engström, B., see Kågedal, L. 537(1991)17

Everaerts, F. M., see Beckers, J. L. 537(1991)407

Everaerts, F. M., see Beckers, J. L. 537(1991)429

Facino, R. M., see Pietta, P. 537(1991)449

Fanska, C. B., Byerley, T. J. and Eick, J. D.
Indirect determination of isocyanates by gas chromatography 537(1991)357

Finn, N., see Smith, R. M. 537(1991)51

Ford, S. H., Gallery, J., Nichols, A. and Shambee, M.
High-performance liquid chromatographic analysis of the (cyanoaquo) stereoisomers of several putative vitamin B_{12} precursors 537(1991)235

Furusawa, M., see Kiba, N. 537(1991)443

Gallery, J., see Ford, S. H. 537(1991)235

Gardana, C., see Pietta, P. 537(1991)449

Gerbino, T. C. and Castello, G.
Gas chromatographic identification of complex mixtures of halomethanes and haloethanes by using the correlation between their retention and vapour pressure 537(1991)305

Goiffon, J.-P., Brun, M. and Bourrier, M.-J.
High-performance liquid chromatography of red fruit anthocyanins 537(1991)101

Goto, J., see Iida, T. 537(1991)345

Goyal, S. S., Hafez, A. and Rains, D. W.
Simultaneous determination of arsenite, arsenate, selenite and selenate in waters using suppressed ion chromatography with ultraviolet absorbance detection 537(1991)269

Guiochon, G., see Czok, M. 537(1991)497

Gülz, P.-G., see Vogt, T. 537(1991)453

Haen, E., Hagenmaier, H. and Remien, J.
Detection of choline and acetylcholine by high-performance liquid chromatography. Limitations, pitfalls, sample preparation 537(1991)514
Hafez, A., see Goyal, S. S. 537(1991)269
Hagenmaier, H., see Haen, E. 537(1991)514
Hamid, H. A., see Al-Ammar, A. S. 537(1991)287
Hasunuma, R., Ogawa, T., Ishii, J. and Kawanishi, Y.
Determination of selenium in organic compounds on a silica gel sintered thin-layer chromatographic plate with 2,3-diaminonaphthalene after direct digestion 537(1991)397
Hermecz, I., see Papp, O. 537(1991)365
Hermecz, I., see Papp, O. 537(1991)371
Hermecz, I., see Papp, O. 537(1991)377
Hina, T., see Kuno, Y. 537(1991)489
Huisden, R. E., Kraak, J. C. and Poppe, H.
Method for the characterization of stationary phases for the separation of proteins by high-performance liquid chromatography 537(1991)1
Hurtubise, R. J., see Mohseni, R. M. 537(1991)61
Iida, T., Komatsubara, I., Chang, F. C., Goto, J. and Nambara, T.
Capillary gas chromatographic behavior of stereoisomeric bile acids with a *vicinal* glycol structure by their "mixed" alkylboronate derivatives 537(1991)345
Ikesaki, T., see Ngan, F. 537(1991)385
Ishii, J., see Hasunuma, R. 537(1991)397
Joelsson, M., see Johansson, G. 537(1991)219
Johansson, G. and Joelsson, M.
Protein–ligand interactions studied on bovine serum albumin with free and polymer-bound Cibacron Blue F3G-A as ligand with reference to affinity partitioning 537(1991)219
Jones, E. A., Bezuidenhout, H. S. and Van Staden, J. F.
Separation of lanthanides and yttrium as anionic complexes by isocratic ion-interaction chromatography 537(1991)277
Kågedal, L., Engström, B., Ellegren, H., Lieber, A.-K., Lundström, H., Sköld, A. and Schenning, M.
Chemical, physical and chromatographic properties of Superdex 75 prep grade and Superdex 200 prep grade gel filtration media 537(1991)17
Kanazawa, H., Nagata, Y., Matsushima, Y., Tomoda, M. and Takai, N.
Preparative high-performance liquid chromatography on chemically modified porous glass. Isolation of acidic saponins from ginseng 537(1991)469
Kaphalia, B. S.
Fatty acid conjugates of chlorinated phenols and their high-performance liquid chromatographic analysis 537(1991)85
Kawanishi, Y., see Hasunuma, R. 537(1991)397
Kiba, N., Shitara, K., Furusawa, M. and Takata, Y.
Post-column oligosaccharide dehydrogenase reactor for coulometric detection of malto-oligosaccharides in a liquid chromatographic system 537(1991)443
Kökösi, J., see Papp, O. 537(1991)365
Kökösi, J., see Papp, O. 537(1991)377
Komatsubara, I., see Iida, T. 537(1991)345
Kraak, J. C., see Huisden, R. E. 537(1991)1
Krull, I. S., see Bourque, A. J. 537(1991)123
Kuno, Y., Hina, T., Akiyama, T. and Matsui, M.
Simultaneous determination of tributyl phosphate and dibutyl phosphate in spent fuel reprocessing streams by gas chromatography 537(1991)489
Kwik, W. L., see Ang, H. G. 537(1991)475
Lanin, S. N. and Nikitin, Y. S.
Molecular interactions in liquid chromatography 537(1991)33
Larionov, O. G., Petrenko, V. V. and Platonova, N. P.
Determination of contributions of different types of solute–sorbent interactions in gas-adsorption chromatography by linear regression of adsorption energies 537(1991)295
Leong, W. K., see Ang, H. G. 537(1991)475
Lieber, A.-K., see Kågedal, L. 537(1991)17
Linde, S., see Welinder, B. S. 537(1991)201
Lundström, H., see Kågedal, L. 537(1991)17
Matsui, M., see Kuno, Y. 537(1991)489
Matsushima, Y., see Kanazawa, H. 537(1991)469
Mauri, P., see Pietta, P. 537(1991)449
Mohseni, R. M. and Hurtubise, R. J.
Changes in the enthalpy and entropy of hydroxyl aromatics in reversed-phase liquid chromatography with β-cyclodextrin in the mobile phase 537(1991)61
Nagata, Y., see Kanazawa, H. 537(1991)469
Nambara, T., see Iida, T. 537(1991)345

Nesterova, I., Rekhter, B., Roshka, G. and Witkiewicz, Z.
Liquid crystals for the gas chromatographic determination of the stereochemistry of insect sex pheromones 537(1991)482
Ngan, F. and Ikesaki, T.
Determination of nine acidic herbicides in water and soil by gas chromatography using an electron-capture detector 537(1991)385
Nichols, A., see Ford, S. H. 537(1991)235
Nikitin, Y. S., see Lanin, S. N. 537(1991)33
Ogawa, T., see Hasunuma, R. 537(1991)397
Olsson, M., Sander, L. C. and Wise, S. A.
Comparison of the liquid chromatographic behaviour of selected steroid isomers using different reversed-phase materials and mobile phase compositions 537(1991)73
Őrfi, L., see Papp, O. 537(1991)371
Pankow, J. F., see Rosen, M. E. 537(1991)321
Papp, O., Szász, G., Kökösi, J. and Hermecz, I.
Correlation between structure and gaschromatographic behaviour of nitrogen-containing heterocyclic compounds. I. Methyl substitution of pyridopyrimidine derivatives 537(1991)365
Papp, O., Szász, G., Kökösi, J. and Hermecz, I.
Correlation between structure and gas chromatographic behaviour of nitrogen-containing heterocyclic compounds. III. Variation of ring size 537(1991)377
Papp, O., Szász, G., Őrfi, L. and Hermecz, I.
Correlation between structure and gas chromatographic behaviour of nitrogen-containing heterocyclic compounds. II. Alkyl substitution of quinazolone derivatives 537(1991)371
Perkins, E. G., see Rojo, J. A. 537(1991)329
Petrenko, V. V., see Larionov, O. G. 537(1991)295
Pietta, P., Mauri, P., Gardana, C., Facino, R. M. and Carini, M.
High-performance liquid chromatographic determination of flavonoid glucosides from *Helichrysum italicum* 537(1991)449
Platonova, N. P., see Larionov, O. G. 537(1991)295
Pontelli, L., see Cartoni, G. P. 537(1991)93
Poppe, H., see Huisden, R. E. 537(1991)1
Quattrucci, E., see Cartoni, G. P. 537(1991)93
Rains, D. W., see Goyal, S. S. 537(1991)269
Rashid, B. H., see Al-Ammar, A. S. 537(1991)287
Rekhter, B., see Nesterova, I. 537(1991)482
Remien, J., see Haen, E. 537(1991)514
Rojo, J. A. and Perkins, E. G.
Isomer identification and gas chromatographic retention studies of monomeric cyclic fatty acid methyl esters 537(1991)329
Rosen, M. E. and Pankow, J. F.
Adsorption/thermal desorption for the determination of volatile organic compounds in water 537(1991)321
Roshka, G., see Nesterova, I. 537(1991)482
Šalamoun, J. and Šlais, K.
Elimination of peak splitting in the liquid chromatography of the proline-containing drug enalapril maleate 537(1991)249
Sander, L. C., see Olsson, M. 537(1991)73
Schenning, M., see Kågedal, L. 537(1991)17
Seidel, A., see Slonina, P. 537(1991)167
Shah, R. S., see Devani, M. B. 537(1991)494
Shah, S. A., see Devani, M. B. 537(1991)494
Shambee, M., see Ford, S. H. 537(1991)235
Shishoo, C. J., see Devani, M. B. 537(1991)494
Shitara, K., see Kiba, N. 537(1991)443
Sigalov, A., Alexandrovich, O. and Strizevskaya, E.
Large-scale isolation and purification of human apolipoproteins A-I and A-II 537(1991)464
Sköld, A., see Kågedal, L. 537(1991)17
Šlais, K., see Šalamoun, J. 537(1991)249
Slonina, P. and Seidel, A.
Breakthrough curves of insulin, [D-Phe^6]-Gonadotropin-releasing hormone and phenylalanine methyl ester on copolymers of alkylacrylate and divinylbenzene 537(1991)167
Smith, R. M. and Finn, N.
Comparison of retention index scales based on alkyl aryl ketones, alkan-2-ones and 1-nitroalkanes for polar drugs on reversed-phase high-performance liquid chromatography 537(1991)51
Snyder, L. R. and Cox, G. B.
Pitfalls in the choice of isotherms for the calculation of band profiles in preparative chromatography. A reply 537(1991)507
Soni, K. P., see Devani, M. B. 537(1991)494
Sørensen, H. H., see Welinder, B. S. 537(1991)181
Speirs, R. J. and Boyer, G. L.
Analysis of ^{55}Fe-labeled hydroxamate siderophores by high-performance liquid chromatography 537(1991)259
Strizevskaya, E., see Sigalov, A. 537(1991)464
Szász, G., see Papp, O. 537(1991)365
Szász, G., see Papp, O. 537(1991)371
Szász, G., see Papp, O. 537(1991)377
Takai, N., see Kanazawa, H. 537(1991)469

Takata, Y., see Kiba, N. 537(1991)443
Thoma, R. S. and Crimmins, D. L.
Automated phenylthiocarbamyl amino acid analysis of carboxypeptidase/aminopeptidase digests and acid hydrolysates 537(1991)153
Tomoda, M., see Kanazawa, H. 537(1991)469
Van Staden, J. F., see Jones, E. A. 537(1991)277
Vogt, T. and Gülz, P.-G.
Isocratic column liquid chromatographic separation of a complex mixture of epicuticular flavonoid aglycones and intracellular flavonol glycosides from *Cistus laurifolius* L. 537(1991)453
Welinder, B. S. and Linde, S.
Reversed-phase high-performance liquid chromatographic characterization of acetic acid extracts of the normal and the diabetic human pancreas 537(1991)201
Welinder, B. S. and Sørensen, H. H.
Alternative mobile phases for the reversed-phase high-performance liquid chromatography of peptides and proteins 537(1991)181
Wise, S. A., see Olsson, M. 537(1991)73
Witkiewicz, Z., see Nesterova, I. 537(1991)482